

Ozurdex Patient Information Leaflet

Ozurdex® has been approved by the National Institute for Health and Care Excellence (NICE) to treat patients with macular oedema (swelling of the back part of the eye) due to uveitis, branch retinal vein occlusion, central retinal vein occlusion or diabetic macular oedema.

How can Ozurdex® help treat macular oedema?

The retina is the light sensitive tissue that lines the back of your eye and is responsible for your eyesight. Macular oedema, a swelling and thickening of the macula, is caused by one of the eye disorders listed above. The macula is a small area in the centre of your retina that contains a rich collection of nerve cells sensitive to light, fine detail, and colour. When the small vessels in the macula leak fluid, this forms a macular oedema or swelling of the retina, as well as causing loss of blood flow and inflammation. All these processes lead to painless reduction of vision in the affected eye. Ozurdex can also be used to treat patients with uveitis (inflammation of the inside of the eye) without macular oedema. Ozurdex® contains dexamethasone, which is slowly released into your eye as the implant

dissolves (biodegrades). Dexamethasone belongs to a group of drugs called potent corticosteroids, and has been shown to reduce inflammation, oedema, and thickening of the retina, therefore improving vision.

Will Ozurdex® treatment fully restore my vision?

Depending on the nature of your eye condition, there is an approximately 50% chance of your vision improving with this treatment. If you decide to undergo treatment, both you and your eye doctor will need to sign and date the informed consent form.

How is Ozurdex administered?

Ozurdex® is a small biodegradable implant which is administered via a special applicator with a needle. It is delivered as an ophthalmic intravitreal injection. This means that it is injected into your eye. Sufficient local anaesthetic is administered prior to the Ozurdex® injection; this procedure is often pain free but not always and you may feel some

discomfort during the injection. The local anaesthetic is given in the form of drops or subconjunctival anaesthetic injection (under the outer covering layer on top of the white of your eye). This injection is usually painless however you may experience brief discomfort.

Are there any side effects associated with Ozurdex®?

All drugs can cause side effects in some people. Some of these side effects may be caused by the injection procedure and not the Ozurdex® implant itself—for example, bleeding on the surface of the eye, difficulties in seeing clearly, a feeling of spots in front of the eye (including 'floaters'), eye pain, seeing flashes of light, swelling or scratching on the surface of the eye, a feeling of looking through mist or fog, redness of the eye, decrease in eye pressure. You may experience temporary visual blurring after receiving an injection and therefore should not drive or use machinery until your vision is recovered. In the days following injection with Ozurdex®, you may experience some of the common side effects listed below. For your information, we have also included the uncommon side effects associated with this treatment, however it is important to remember that these are rare.

Very common side effects - might affect more than 1 patient in 10:

- Increased pressure in the eye, bleeding on the surface of the eye.

Common side effects - might affect 1 to 10 patients in 100:

- High pressure in the eye. Separation of the jelly inside the eye (vitreous detachment). Clouding of the eye lens (cataract). Bleeding inside the eye. Difficulties in seeing clearly. A feeling of spots in front of the eye (including 'floaters'). Eye pain. Seeing flashes of light. Swelling or scratching on the surface of the eye. A feeling of looking through mist or fog. Redness of the eye

Uncommon side effects - might affect one to 10 patients in 1,000:

- Retinal tear. Inflammation. Decrease in eye pressure, infection inside the eye, severe bleeding inside the eye

Aftercare advice:

You can continue normal activities once you have been discharged from hospital. Please use the post-procedure antibiotic drops as advised. If you experience eye pain or loss of vision, please contact us without delay. You should also seek immediate care if your eye becomes red, sensitive to light, painful, or develops a change in vision.

It is important to attend your post procedure checks to monitor the pressure inside your eyes. If the eye pressure is too high for too long and is not treated, there can be damage to your vision.

A follow up appointment will be sent to you for your intra ocular pressure to be checked in 3-4 weeks following your Ozurdex implant procedure.

How to contact the eye department after surgery

For post-surgery queries or problems,
Monday – Saturday between 09:00 and
17:00

Call our Eye Triage Team 01392 404955

Alternatively, you can leave a **call back request** with the Day Case staff on
01932 406013.

Monday to Friday between 07:30 and
17:00

Outside normal eye department working hours

Please call **111** for emergencies and
urgent queries only.

The Trust cannot accept any responsibility for the accuracy of the information given if the leaflet is not used by Royal Devon staff undertaking procedures at the Royal Devon hospitals.

© Royal Devon University Healthcare NHS Foundation Trust

Designed by Graphics (Print & Design), RD&E (Heavitree)